arXiv:hep-th/9904103v3 10 Nov 1999

# FORMULAS FOR $A_n$ AND $B_n$–SOLUTIONS OF WDVV EQUATIONS

S.. NATANZON

## 1. Introduction

1. WDVV equations were appeared in works of E.Witten [W1], R.Dijkgraaf, E.Verlinde, H.Verlinde [DVV] as equations for primary free energy of two-dimensional topological field theories. The simplest non-trivial solutions of WDVV equations are $A_n$ and $B_n$-potentials, which describe also metrics of K.Saito on spaces of versal deformation of $A_n$ and $B_n$-singularities. According to E.Witten [W2] coefficients of $A_n$–potentials form intersection numbers of Mumford–Morita–Miller classes of a moduli space of spheres with punctures and $n$–spin structures. The potentials are known for $n \leqslant 4$ [W2, D1, D2]. We find these potentials for all $n$, using its connection with Gelfand-Dikii $(n - k\partial V)$ hierarchy. In passing we give recurrence formulas for coefficients of dispersionless KP hierarchy. Results of J.-B.Zuber [Z] give a reduction of $B_n$–potentials to $A_n$–potentials.

I would like to thank B.Dubrovin for fruitful discussions.

## 2. Main theorems

For calculation of the $A_{n-1}$-potential we use combinatorial coefficients $P_n(i_1 ... i_m)$, where $n$ and $s_j$ are natural numbers. Put us

$$P_n(i) = n, \quad P_n(s_1 ... s_m) = C_n^m - \sum_{q=1}^{m-1} P_n(s_1 ... s_q) C^{m-q}_{n-q-(s_1+\cdots+s_q)},$$

where here and later

$$0! = 1, C_p^t = \frac{p!}{t!(p-t)!} \text{ for } p \geqslant t \geqslant 0 \text{ and } C_p^t = 0 \text{ in another cases.}$$

Put us $x_0 = 0$. For $q > 0$ we recurrently define polynomials $x_{-q}$ $(x_1, ..., x_{n-1})$ putting

$$x_{-q} = -\frac{1}{n} \sum_{m=2}^{\infty} \sum P_n(s_1 \ldots s_m) x_{n-s_1} \cdots x_{n-s_m},$$

where the second sum is taken by all natural numbers $(s_1, ..., s_m)$ such that $\sum_{i=1}^m s_i = q+n+1-m$.

This polynomial has a form

$$x_{-q} = \sum_{m=1}^{\infty} \sum B_{qi_1 \cdots i_m} x_{i_1} \cdots x_{i_m},$$

where $B_{qi_1,...,i_m}$ are constants and the second sum is taken by numbers $(i_1, ..., i_m)$ such that $1 \leqslant i_1 \leqslant ... \leqslant i_m \leqslant n-1$.

Put us

$$A_{i_1 \cdots i_m} = -\frac{1}{i_1 p} B_{i_1 \cdots i_m}$$

and $p$ is the number of $j$ such that $i_j = i_1$.

Consider

$$F_A = \sum_{m=3}^{\infty} \sum A_{i_1 \cdots i_m} x_{i_1} \cdots x_{i_m},$$

$$F_B = \sum_{m=3}^{\infty} \sum A_{2i_1-1 \cdots 2i_m-1} x_{i_1} \cdots x_{i_m},$$

where the second sums is taken by $1 \leqslant i_1 \leqslant ... \leqslant i_m \leqslant n-1$.

**Theorem 1.** *Polynomial $F_A$ is a $A_{n-1}$–potential. Polynomial $F_B$ is a $B_{n-1}$–potential. This means that they satisfy the equations*

$$\sum_{\gamma=1}^{n-1} \frac{\partial^3 F}{\partial x_\alpha \partial x_\beta \partial x_\gamma} \frac{\partial^3 F}{\partial x_{n-\gamma} \partial x_\eta \partial x_\xi} = \sum_{\gamma=1}^{n-1} \frac{\partial^3 F}{\partial x_\xi \partial x_\beta \partial x_\gamma} \frac{\partial^3 F}{\partial x_{n-\gamma} \partial x_\eta \partial x_\alpha}, \ (\alpha, \beta, \eta, \xi = 1, ..., n-1),$$

$$\frac{\partial^3 F}{\partial x_1 \partial x_\alpha \partial x_\beta} = \delta_{\alpha+\beta,n}, \quad \sum_{j=1}^{n-1} d_i \frac{\partial F}{\partial x_i} = dF,$$

*where $d_i = 1 + \frac{1-i}{n}$, $d = 2 + \frac{2}{n}$ for $F = F_A$ and $d_i = 1 + \frac{1-i}{n-1}$, $d = 2 + \frac{1}{n-1}$ for $F = F_B$.*

## 3. Combinatorial lemma

For natural $s, i_1, ..., i_n$ and integer not negative $j_1, ..., j_n$ defind $P_s\begin{pmatrix} i_1 & \cdots & i_n \\ j_1 & \cdots & j_n \end{pmatrix}$ by recurence formulas:

$$1) P_s\begin{pmatrix} i_1 & \cdots & i_n \\ 0 & \cdots & 0 \end{pmatrix} = 0; \quad 2) P_s\begin{pmatrix} i \\ j \end{pmatrix} = C_s^j \quad \text{for } j > 0;$$

$$3)P_s\begin{pmatrix} i_1 & \dots & i_n \\ j_1 & \dots & j_n \end{pmatrix} = \frac{1}{n!}C_s^{j_1+\dots+j_n}\frac{(j_1+\dots+j_n)!}{j_1!\cdots j_n!} -$$

$$\sum_{q=1}^{n-1} P_s\begin{pmatrix} i_1 & \dots & i_q \\ j_1 & \dots & j_q \end{pmatrix}\frac{1}{(n-q)!}C_{s-(i_1+\dots+i_q+j_1+\dots+j_q)}^{j_{q+1}+\dots+j_n}\frac{(j_{q+1}+\dots+j_n)!}{j_{q+1}!\cdots j_n!}$$

for $(j_1, ..., j_n) \neq (0, ..., 0)$.

Let $\begin{bmatrix} i_1 & \dots & i_n \\ j_1 & \dots & j_n \end{bmatrix}$ be the set of all matrices, which appear from $\begin{pmatrix} i_1 & \dots & i_n \\ j_1 & \dots & j_n \end{pmatrix}$ by permutation of columns. Let $\begin{Vmatrix} i_1 & \dots & i_n \\ j_1 & \dots & j_n \end{Vmatrix}$ be the number of such matrices. Put us

$$P_s\begin{bmatrix} i_1 & \dots & i_n \\ j_1 & \dots & j_n \end{bmatrix} = \sum P_s\begin{pmatrix} a_1 & \dots & a_n \\ b_1 & \dots & b_n \end{pmatrix},$$

where the sum is taken by all

$$\begin{pmatrix} a_1 & \dots & a_n \\ b_1 & \dots & b_n \end{pmatrix} \in \begin{bmatrix} i_1 & \dots & i_n \\ j_1 & \dots & j_n \end{bmatrix}$$

**Lemma 1** *Let $m > 0, k > 0$ and $j_n \geqslant 1$ for $n \leqslant m$. Then*

$$P_s\begin{bmatrix} i_1 & \dots & i_m & i_{m+1} & \dots & i_{m+k} \\ j_1 & \dots & j_m & 0 & \dots & 0 \end{bmatrix} =$$

$$= \begin{cases} 0, & \textit{if } s \geqslant i_1+...+i_m+j_1+...+j_m, \\ \frac{1}{k!}\begin{Vmatrix} i_{m+1} & \dots & i_{m+k} \\ 0 & \dots & 0 \end{Vmatrix} P_s\begin{bmatrix} i_1 & \dots & i_m \\ j_1 & \dots & j_m \end{bmatrix}, & \textit{if } s < i_1+...+i_m+j_1+...+j_m. \end{cases}$$

Proof: Prove at first the lemma for $m = 1$ using induction by k. For $m = k = 1$

$$P_s\begin{bmatrix} i_1 & i_2 \\ j_1 & 0 \end{bmatrix} = P_s\begin{pmatrix} i_1 & i_2 \\ j_1 & 0 \end{pmatrix} + P_s\begin{pmatrix} i_2 & i_1 \\ 0 & j_1 \end{pmatrix} =$$

$$\frac{1}{2}C_s^{j_1} - P_s\begin{pmatrix} i_1 \\ j_1 \end{pmatrix}\cdot C_{s-(i_1+j_1)}^0 + \frac{1}{2}C_s^{j_1} - P_s\begin{pmatrix} i_2 \\ 0 \end{pmatrix}\cdot C_{s-i_2}^0 = C_s^{j_1} - C_s^{j_1}C_{s-(i_1+j_1)}^0 =$$

$$= \begin{cases} 0, & \text{if } s \geqslant i_1+j_1, \\ C_s^{j_1} = P_s\begin{bmatrix} i_1 \\ j_1 \end{bmatrix}, & \text{if } s < i_1+j_1. \end{cases}$$

Prove the lemma for $m = 1, k = N$, considering that it is proved for $m = 1, k < N$. If $s \geqslant i_1 + j_1$, then

$$P_1\begin{bmatrix} i_1 & i_2 & \dots & i_{k+1} \\ j_1 & 0 & \dots & 0 \end{bmatrix} = \frac{1}{k!}C_s^{j_1}\left\|\begin{matrix} i_2 & \dots & i_{k+1} \\ 0 & \dots & 0 \end{matrix}\right\| - P_s\begin{pmatrix} i_1 \\ j_1 \end{pmatrix}\cdot C^0_{s-(i_1+j_1)}\cdot\frac{1}{k!}\left\|\begin{matrix} i_2 & \dots & i_{k+1} \\ 0 & \dots & 0 \end{matrix}\right\| = 0.$$

If $s < i_1 + j_1$ than

$$P_1\begin{bmatrix} i_1 & i_2 & \dots & i_{k+1} \\ j_1 & 0 & \dots & 0 \end{bmatrix} = \frac{1}{k!}C_s^{j_1}\left\|\begin{matrix} i_2 & \dots & i_{k+1} \\ 0 & \dots & 0 \end{matrix}\right\| - A\cdot C^0_{s-(i_1+j_1)} =$$

$$\frac{1}{k!}\left\|\begin{matrix} i_2 & \dots & , i_{k+1} \\ 0 & \dots & 0 \end{matrix}\right\| P_s\begin{bmatrix} i_1 \\ j_1 \end{bmatrix}.$$

Thus the lemma is proved for $m = 1$.

Prove the lemma for $m = N$ considering that it is proved for $m < N$. Then

$$P_s\begin{bmatrix} i_1 & \dots & i_m & i_{m+1} & \dots & i_{m+k} \\ j_1 & \dots & j_m & 0 & \dots & 0 \end{bmatrix} =$$

$$\sum_{\begin{pmatrix} \alpha_1 & \dots & \alpha_m \\ \beta_1 & \dots & \beta_m \end{pmatrix} \in \begin{bmatrix} i_1 & \dots & i_m \\ j_1 & \dots & j_m \end{bmatrix}} \Bigg(\frac{1}{(m+k)!}C_s^{\beta_1+\dots+\beta_m}\frac{(\beta_1+\dots+\beta_m)!}{\beta_1!\cdots\beta_m!}C^k_{m+k}\cdot$$

$$\left\|\begin{matrix} i_{m+1} & \dots & i_{m+k} \\ 0 & \dots & 0 \end{matrix}\right\| -$$

$$-\sum_{q=1}^{m} P_s\begin{pmatrix} \alpha_1 & \dots & \alpha_q \\ \beta_1 & \dots & \beta_q \end{pmatrix}\frac{1}{(m+k-q)!}C^{\beta_{q+1}+\dots+\beta_m}_{s-(\alpha_1+\dots+\alpha_q+\beta_1+\dots+\beta_q)}\frac{(\beta_{q+1}+\dots+\beta_m)!}{\beta_{q+1}!\cdots\beta_m!}C^k_{m+k-q}\cdot$$

$$\left\|\begin{matrix} i_{m+1} & \dots & i_{m+k} \\ 0 & \dots & 0 \end{matrix}\right\|\Bigg) = P_s\begin{bmatrix} i_1 & \dots & i_m \\ j_1 & \dots & j_m \end{bmatrix}\frac{1}{k!}\left\|\begin{matrix} i_{m+1} & \dots & i_{m+k} \\ 0 & \dots & 0 \end{matrix}\right\| -$$

$$P_s\begin{bmatrix} i_1 & \dots & i_m \\ j_1 & \dots & j_m \end{bmatrix} C^0_{s-(i_1+\dots+i_m+j_1+\dots+j_m)}\frac{1}{k!}\left\|\begin{matrix} i_{m+1} & \dots & i_{m+k} \\ 0 & \dots & 0 \end{matrix}\right\| =$$

$$= \begin{cases} 0, & \text{if } s \geqslant i_1+\dots+i_m+j_1+\dots+j_m, \\ P_s\begin{bmatrix} i_1 & \dots & i_m \\ j_1 & \dots & j_m \end{bmatrix}\frac{1}{k!}\left\|\begin{matrix} i_{m+1} & \dots & i_{m+k} \\ 0 & \dots & 0 \end{matrix}\right\|, & \text{if } s < i_1+\dots+i_m+j_1+\dots+j_m. \square \end{cases}$$

## 4. Equations for Bacher-Akhiezer function

Consider the KP hierarchy. This is a condition of compatibility of the infinite system of the differential equations

$$\frac{\partial\psi}{\partial x_n} = L_n\psi, \tag{1}$$

where

$$L_n = \frac{\partial^n}{\partial x_1^n} + \sum_{i=2}^{n} B_n^i(x)\frac{\partial^{n-i}}{\partial^{n-i}x_1},$$

and $\psi$ is a function of type

$$\psi(x,k) = \exp(\sum_{j=1}^{\infty} x_j k^j)(1+\sum_{i=1}^{\infty}\xi_i k^{-i}),$$

(here $k \in \mathbb{C}$ belong to some neighbourhood of $\infty$ and $x = (x_1, x_2, ...)$ — is a finite sequence).

Put us

$$\partial_i = \frac{\partial}{\partial x_i},\ \partial = \partial_1.$$

A direct calculation gives

**Lemma 2.** *Conditions of compatibility of (1) are*

$$B_s^t = -\sum_{i=1}^{t-1} C_s^i \partial^i \xi_{t-i} - \sum_{j=2}^{t-1} B_s^j \sum_{i=0}^{t-j-1} C_{s-j}^i \partial^i \xi_{t-i-j}, \tag{2}$$

$$\partial_n\xi_i = \sum_{j=1}^{n+i-1} C_n^j \partial^j \xi_{i+n-j} + \sum_{k=2}^{n} B_n^k \sum_{j=0}^{n-k} C_{n-k}^j \partial^j \xi_{i+n-j-k}. \quad \Box \tag{3}$$

In this case $\psi$ is called a *Bacher-Akhiezer function.*

Consider now the function

$$\ln\psi(x,k) = \sum_{j=1}^{\infty} x_j k^j + \sum_{j=1}^{\infty}\eta_j k^{-j},$$

where

$$\xi_j = \sum_{n=1}^{\infty}\frac{1}{n!}\sum_{i_1+\cdots+i_n=j}\eta_{i_1}\cdots\eta_{i_n}.$$

**Lemma 3.** *Let $2 \leqslant t \leqslant s$. Then*

$$B_s^t = -\sum_{n=1}^{\infty} \sum P_s \begin{pmatrix} i_1 & \dots & i_n \\ j_1 & \dots & j_n \end{pmatrix} \partial^{j_1} \eta_{i_1} \cdots \partial^{i_n} \eta_{i_n},$$

*where the second sum is taken by all matrices* $\begin{pmatrix} i_1 & \dots & i_n \\ j_1 & \dots & j_n \end{pmatrix}$ *such that* $i_m \geqslant 1, j_m \geqslant 1$ *and* $i_1 + \cdots + i_n + j_1 + \cdots + j_n = t$.

Proof: An induction by $t$. For $t = 2$ according to (2), $B_s^2 = -s\partial\xi_1 = -P_s\binom{1}{1}\partial\eta_1$. Prove the lemma for $t = N$ considering that it is proved for $t < N$. According to (2)

$$B_s^t = -\sum_{i=1}^{t-1} C_s^i \partial^i \Big( \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{i_1+\cdots+i_n=t-i} \eta_{i_1} \cdots \eta_{i_n} \Big) +$$

$$+ \sum_{j=2}^{t-1} \Big( \sum_{n=1}^{\infty} \sum_{i_1+\cdots+j_n=j} P_s \begin{pmatrix} i_1 & \dots & i_n \\ j_1 & \dots & j_n \end{pmatrix} \partial^{j_1} \eta_{i_1} \cdots \partial^{j_n} \eta_{i_n} \Big) \cdot$$

$$\Big( \sum_{i=0}^{t-j-1} C_{s-j}^i \partial^j \Big( \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{i_1+\cdots+i_n=t-i-j} \eta_{i_1} \cdots \eta_{i_n} \Big) \Big) =$$

$$= -\sum_{n=1}^{\infty} \sum \Big( \frac{1}{n!} C_s^{j_1+\cdots+j_n} \frac{(j_1 + \cdots + j_n)!}{j_1! \cdots j_n!} -$$

$$\sum_{q=1}^{n-1} P_s \begin{pmatrix} i_1 & \dots & i_q \\ j_1 & \dots & j_q \end{pmatrix} \frac{1}{(n-q)!} C_{s-(i_1+\cdots i_q+j_1+\cdots+j_q)}^{j_{q+1}\cdots j_n} \frac{(j_{q+1} + \cdots j_n)!}{j_{q+1}! \cdots j_n!} \Big) \partial^{j_1} \eta_{i_1} \cdots \partial^{j_n} \eta_{i_n} =$$

$$= -\sum_{n=1}^{\infty} \sum P_s \begin{pmatrix} i_1 & \dots & i_n \\ j_1 & \dots & j_n \end{pmatrix} \partial^{j_1} \eta_{i_1} \cdots \partial^{j_n} \eta_{i_n},$$

where the second sums are taken by all matrices $\begin{pmatrix} i_1 & \dots & i_n \\ j_1 & \dots & j_n \end{pmatrix}$ such that $i_1 + \cdots + i_n + j_1 + \cdots + j_n = t$, $i_m \geqslant 1, j_m \geqslant 0$. According to lemma 1 it is possible to consider that in the last sum $j_m > 0$ for all $m$. □

**Lemma 4.**

$$\partial_s \eta_r = \sum_{n=1}^{\infty} \sum P_s \begin{pmatrix} i_1 & \dots & i_n \\ j_1 & \dots & j_n \end{pmatrix} \partial^{j_1} \eta_{i_1} \cdots \partial^{j_n} \eta_{i_n},$$

*where the second sum is taken by all matrices* $\begin{pmatrix} i_1 & \cdots & i_n \\ j_1 & \cdots & j_n \end{pmatrix}$ *such that* $i_m \geqslant 1, j_m \geqslant 1$ *and* $i_1 + \cdots + i_n + j_1 + \cdots + j_n = r + s$.

Proof: An induction by $r$. According to (3) and lemma 3 for $r = 1$

$$\partial_s \eta_1 = \partial_s \xi_1 = \sum_{j=1}^{\infty} C_s^j \partial^j \xi_{s+1-j} + \sum_{k=2}^{s} B_s^k \sum_{j=0}^{s-k} C_{s-k}^j \partial^j \xi_{1+s-j-k} =$$

$$\sum_{j=1}^{s} C_s^j \partial^j \xi_{s+1-j} - \sum_{k=2}^{\infty} \Big( \sum_{n=1}^{\infty} \sum_{i_1+\cdots+j_n=k} P_s \begin{pmatrix} i_1 & \cdots & i_n \\ j_1 & \cdots & j_n \end{pmatrix} \partial^{j_1} \eta_{i_1} \cdots \partial^{j_n} \eta_{i_n} \Big) \cdot$$

$$\Big( \sum_{j=0}^{s-k} C_{s-k}^j \partial^j \xi_{1+s-j-k} \Big) = \sum_{n=1}^{\infty} \sum P_s \begin{pmatrix} i_1 & \cdots & i_n \\ j_1 & \cdots & j_n \end{pmatrix} \partial^{j_1} \eta_{i_1} \cdots \partial^{j_n} \eta_{i_n},$$

where the second sum in the last formula is taken by all matrices $\begin{pmatrix} i_1 & \cdots & i_n \\ j_1 & \cdots & j_n \end{pmatrix}$ such that $i_1 + \cdots + i_n + j_1 + \cdots + j_n = s + 1,\ i_m \geqslant 1,\ j_m \geqslant 0$. According to lemma 1 in this sum it is sufficient consider only matrices, where $j_m > 0$ for all $m$.

Prove now the lemma for $r = N$, considering that it is proved for $r < N$. According to (3)

$$\partial_s \Big( \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{i_1+\cdots+i_n=r} \eta_{i_1} \cdots \eta_{i_n} \Big) = \sum_{j=1}^{s+r-1} C_s^j \partial^j \xi_{s+r-j} + \sum_{k=2}^{\infty} B_s^k \sum_{j=0}^{s-k} C_{s-k}^j \partial^j \xi_{r+s-j-k}.$$

Thus according to lemma 3, lemma 1 and inductive hypothesis,

$$\partial_s \eta_r = \sum_{j=1}^{\infty} C_s^j \partial^j \Big( \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{i_1+\cdots+i_n=s+r-j} \eta_{i_1} \cdots \eta_{i_n} \Big) +$$

$$+ \sum_{k=2}^{\infty} \Big( \sum_{i_1+\cdots+j_n=k} P_s \begin{pmatrix} i_1 & \cdots & i_n \\ j_1 & \cdots & j_n \end{pmatrix} \partial^{j_1} \eta_{i_1} \cdots \partial^{j_n} \eta_{i_n} \Big) \cdot$$

$$\sum_{j=0}^{s-k} C_{s-k}^j \partial^j \Big( \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{i_1+\cdots+i_n=r+s-j-k} \eta_{i_1} \cdots \eta_{i_n} \Big) - \partial_s \Big( \sum_{n=2}^{\infty} \frac{1}{n!} \sum_{i_1+\cdots+i_n=r} \eta_{i_1} \cdots \eta_{i_n} \Big) =$$

$$= \sum_{n=1}^{\infty} \sum P_s \begin{pmatrix} i_1 & \cdots & i_n \\ j_1 & \cdots & j_n \end{pmatrix} \partial^{j_1} \eta_{i_1} \cdots \partial^{j_n} \eta_{i_n},$$

where the second sum in the last formula is taken by all matrices $\begin{pmatrix} i_1 & \dots & i_n \\ j_1 & \dots & j_n \end{pmatrix}$ such that $i_1 + \dots + i_n + j_1 + \dots + j_n = s + r$, $i_m \geqslant 1, j_m \geqslant 1$. □

## 5. KP hierarchy

According to [DKJM] the Bacher-Akhiezer function $\psi$ is

$$\psi(x,k) = \exp(\sum x_j k^j) \frac{\tau(x_1 - k^{-1}, x_2 - \frac{1}{2}k^{-2}, x_3 - \frac{1}{3}k^{-3}, ...)}{\tau(x_1, x_2, x_3, ...)}$$

for some function $\tau(x_1, x_2, ...)$. By analogy of [N] this gives a possibility to describe the KP hierarchy as an infinite system of differential equations on $v(x,k) = -\ln \tau(x,k)$. Really

$$\sum_{j=1}^{\infty} \eta_j k^{-j} = \ln \psi(x,k) - \sum_{j=1}^{\infty} x_j k^j = -v(x_1 - k^{-1}, x_2 - \frac{1}{2}k^{-2}, ...) + v(x) =$$

$$\sum_{n=1}^{\infty} \sum_{i_1 + \dots + i_n = j} \frac{(-1)^{n+1}}{n! i_1 \dots i_n} \partial_{i_1} \cdots \partial_{i_n} v(x) k^{-j}.$$

Therefore

$$\eta_r = \sum_{n=1}^{\infty} \sum_{i_1 + \dots + i_n = r} \frac{(-1)^{n+1}}{n! i_1 \dots i_n} \partial_{i_1} \cdots \partial_{i_n} v. \tag{4}$$

**Theorem 2.** *There exist universal rational coefficients*

$$R_r \begin{pmatrix} s_1 & \dots & s_n \\ t_1 & \dots & t_n \end{pmatrix}, R_{ij} \begin{pmatrix} s_1 & \dots & s_n \\ t_1 & \dots & t_n \end{pmatrix}$$

*such that*

$$\eta_r = \frac{1}{r} \partial_r v + \sum_{n=1}^{\infty} \sum R_r \begin{pmatrix} s_1 & \dots & s_n \\ t_1 & \dots & t_n \end{pmatrix} \partial_{s_1} \partial^{t_1} v \cdots \partial_{s_n} \partial^{t_n} v, \tag{5}$$

$$\partial_i \partial_j v = \sum_{n=1}^{\infty} \sum R_{ij} \begin{pmatrix} s_1 & \dots & s_n \\ t_1 & \dots & t_n \end{pmatrix} \partial_{s_1} \partial^{t_1} v \cdots \partial_{s_n} \partial^{t_n} v, \tag{6}$$

*where the second sums are taken by all matrices* $\begin{pmatrix} s_1 & \dots & s_n \\ t_1 & \dots & t_n \end{pmatrix}$ *such that* $s_m, t_m \geqslant 1$, *and the sum* $s_1 + \dots + s_n + t_1 + \dots + t_n$ *is equal* $r$ *for (5) and* $i + j$ *for (6). Moreover*

$$R_{ij} \begin{pmatrix} s_1 & \dots & s_n \\ 1 & \dots & 1 \end{pmatrix} = \frac{j}{s_1 \cdots s_n} P_i(s_1 ... s_n).$$

Proof: An induction by $k$ and $i+j$. For $i+j=2$ the theorem is obviously. For $r=1$ it follows from (4). Prove the theorem for $i+j=N$ and $r=N-1$, considering that it is proved for $i+j<N$ and $r<N-1$. Later we consider that $s_m, t_m \geqslant 1$ and $\sigma_n = s_1+\cdots+s_n+t_1+\cdots+t_n$. Then according to (4) and (6)

$$\eta_r = \frac{1}{r}\partial_r v + \sum_{n=2}^{\infty} \sum_{s_1+\cdots+s_n=r} \frac{(-1)^{n+1}}{n! s_1\cdots s_n}\partial_{s_1}\cdots\partial_{s_n} v(x) =$$

$$\frac{1}{r}\partial_r v + \sum_{n=1}^{\infty}\sum_{\sigma_n=r} R_r \begin{pmatrix} s_1 & \dots & s_n \\ t_1 & \dots & t_n \end{pmatrix} \partial_{s_1}\partial^{t_1} v \cdots \partial_{s_n}\partial^{t_n} v.$$

Thus according to (5), (6) and lemma 4

$$\frac{1}{j}\partial_i\partial_j v = \partial_i\eta_j - \partial_i\Big(\sum_{n=1}^{\infty}\sum_{\sigma_n=j} R_j \begin{pmatrix} s_1 & \dots & s_n \\ t_1 & \dots & t_n \end{pmatrix} \partial_{s_1}\partial^{t_1} v \cdots \partial_{s_n}\partial^{t_n} v\Big) =$$

$$\sum_{n=1}^{\infty}\sum_{\sigma_n=i+j} P_i \begin{pmatrix} s_1 & \dots & s_n \\ t_1 & \dots & t_n \end{pmatrix} \partial^{t_1}\eta_{s_1}\cdots\partial^{t_n}\eta_{s_n} -$$

$$\partial_i\Big(\sum_{n=1}^{\infty}\sum_{\sigma_n=j} R_j \begin{pmatrix} s_1 & \dots & s_n \\ t_1 & \dots & t_n \end{pmatrix} \partial^{t_1}\partial_{s_1} v \cdots \partial^{t_n}\partial_{s_n} v\Big) =$$

$$\sum_{n=1}^{\infty}\sum_{s_1+\cdots+s_n+n=i+j} P_i \begin{pmatrix} s_1 & \dots & s_n \\ 1 & \dots & 1 \end{pmatrix} \partial(\frac{1}{s_1}\partial_{s_1} v)\cdots\partial(\frac{1}{s_n}\partial_{s_n} v) +$$

$$\sum_{n=1}^{\infty}\sum_{\sigma_n=i+j, t_1+\cdots+t_n>n} R_{ij} \begin{pmatrix} s_1 & \dots & s_n \\ t_1 & \dots & t_n \end{pmatrix} \partial_{s_1}\partial^{t_1} v \cdots \partial_{s_n}\partial^{t_n} v.$$

The obvious relation $P_i \begin{pmatrix} s_1 & \dots & s_n \\ 1 & \dots & 1 \end{pmatrix} = P_i(s_1 ... s_n)$ conclude the proof. □

**Remark.** The first equation from (6) is the "integrated over $x_1$" KP equation

$$\partial_2^2 v = \frac{4}{3}\partial_3\partial_1 v - \frac{1}{3}\partial_1^4 v + 2(\partial_1^2 v)^2.$$

The relation (6) was first deduced in [ DN ] from the Hirota hierarchy [ DKJM ]. This method gives another formulas for coefficients $R_{ij}$. The equalities between the coefficients provide nontrivial combinatorial identities. □

The system (6) is called the *KP hierarchy. The dispersionless KP hierarchy* is called the infinite system of differential equations which appears from (6) if to throw out all monomials

$$R_{ij}\begin{pmatrix} s_1 & \dots & s_n \\ t_1 & \dots & t_n \end{pmatrix} \partial_{s_1}\partial^{t_1}v\cdots\partial_{s_n}\partial^{t_n}v,$$

where $\sum_{i=1}^{n} t_j > n$. The theorem 2 gives

**Consequence** *The despersionless KP hierarchy is*

$$\frac{1}{j}\partial_i\partial_j v = \sum_{m=1}^{\infty}\sum_{s_1+\dots+s_m=i+j-m} P_i(s_1 ... s_m)\frac{1}{s_1}\partial\partial_{s_1}v\cdots\frac{1}{s_m}\partial\partial_{s_m}v. \qquad (7)$$

## 6. Proof of theorem 1

According to [DVV, K, W2] one of $A_{n-1}$-potentials is

$$\widetilde{F}(x_1, ..., x_{n-1}) = v(x_1, \frac{x_2}{2}, ..., \frac{x_{n-1}}{n-1}, 0, ...),$$

where $v$ satisfies the dispersionless KP hierarchy and $\partial_n v = 0$ (dispersionless Gelfand-Dikii equation).

According to (7) from this follows

$$0 = \partial_\ell\partial_n\widetilde{F} = \sum_{m=1}^{\infty}\sum_{s_1+\dots+s_m=\ell+n-m} P_n(s_1\dots s_m)\partial\partial_{s_1}\widetilde{F}\cdots\partial\partial_{s_m}\widetilde{F} =$$

$$n\partial\partial_{n+\ell-1}\widetilde{F} + \sum_{m=2}^{\infty}\sum_{s_1+\dots+s_m=\ell+n-m} P_n(s_1\dots s_m)\partial\partial_{s_1}\widetilde{F}\cdots\partial\partial_{s_m}\widetilde{F}.$$

Thus

$$\partial\partial_{n+\ell-1}\widetilde{F} = -\frac{1}{n}\sum_{m=2}^{\infty}\sum_{s_1+\dots+s_m=\ell+n-m} P_n(s_1\dots s_m)\partial\partial_{s_1}\widetilde{F}\cdots\partial\partial_{s_m}\widetilde{F}. \qquad (8)$$

According to [W2]

$$\partial\partial_s\widetilde{F} = x_{n-s} \quad \text{and} \quad \partial\partial_{n+s}\widetilde{F} = -s\partial_s\widetilde{F}$$

for $s < n$. In additional $\partial\partial_n\widetilde{F} = 0$ and thus according to (8) $\partial\partial_{n+\ell-1}\widetilde{F} = x_{1-\ell}$ for all $\ell$. Therefore

$$\partial_s\widetilde{F} = -\frac{1}{s}x_{-s} = \partial_s F_A$$

and $F_A - \widetilde{F}$ — is a linear form. According to [D2, p.16] this form is equal 0. According [Z] $F_B(x_1, ..., x_n) = F_A(x_1, 0, x_2, 0, ..., x_{n-1}, 0, x_n)$.□

There are the potentials finding by our formulas

$$(A_2) \quad F = \frac{1}{2}x_1^2x_2 + \frac{1}{24}x_2^4,$$

$$(A_3) \quad F = \frac{1}{2}x_1^2x_3 + \frac{1}{2}x_1x_2^2 + \frac{1}{4}x_2^2x_3^2 + \frac{1}{60}x_3^5,$$

$$(A_4) \quad F = \frac{1}{2}x_1^2x_4 + x_1x_2x_3 + \frac{1}{6}x_2^3 + \frac{1}{12}x_3^4 + \frac{1}{2}x_2x_3^2x_4 + \frac{1}{4}x_2^2x_4^2 + \frac{1}{6}x_3^2x_4^3 + \frac{1}{120}x_4^6,$$

$$(A_5) \quad F = \frac{1}{2}x_1^2x_5 + \frac{1}{2}x_2^2x_3 + \frac{1}{4}x_2^2x_5^2 + x_2x_3x_4x_5 + x_1x_2x_4 + \frac{1}{6}x_2x_4^3 + \frac{1}{2}x_1x_3^2 + \frac{1}{2}x_3^2x_4^2 +$$

$$+\frac{1}{6}x_3^3x_5 + \frac{1}{6}x_3^2x_5^3 + \frac{1}{2}x_3x_4^2x_5^2 + \frac{1}{6}x_4^4x_5 + \frac{1}{8}x_4^2x_5^4 + \frac{1}{210}x_5^7.$$

$$(B_2) \quad F = \frac{1}{2}x_1^2x_2 + \frac{1}{60}x_2^5,$$

$$(B_3) \quad F = \frac{1}{2}x_1^2x_3 + \frac{1}{2}x_1x_2^2 + \frac{1}{6}x_2^3x_3 + \frac{1}{6}x_2^2x_3^3 + \frac{1}{210}x_3^7.$$

## References

[D1] B.Dubrovin, Geometry of 2D topological field theories, In: "Integrable Systems ans Quantum Groups", Eds. M.Francaviglia, S.Greco,Springer Lecture Notes in Math. 1620 (1996), 120-348.

[D2] B.Dubrovin, Painleve transcendents in two-dimensional topological field theory. Preprint SISSA 24/98/FM P.107.

[DKJM] E.Date, M.Kashiwara, M.Jimbo, T.Miwa, Transformation groups for soliton equation, Proceedings of RIMS Symposium on Non-Linear Integrable System. Singapore: World Science Publ. Co., 1983, 39-119.

[DN] B.A.Dubrovin, S.M.Natanzon, Real theta-function solutions of the Kadomtsev-Petviashvili equation. Math. USSR Irvestiya, 32:2 (1989), 269-288.

[DVV] R.Dijkgraaf, E.Verlinde, H.Verlinde, Topological strings in $d < 1$. Nucl. Phys., B 352 (1991), 59.

[K] I.Krichever, The dispersionless Lax equations and topological minimal models. Commun. Math. Phys. 143 (1992), 415-429.

[N] S.M.Natanzon, Differentil equations on the Prym theta function. A realness criterion for two-dimensional, finite-zone, potential Schrödinger operators. Functional Anal. Appl. 26:1 (1992), 13-20.

[W1] E.Witten, On the structure of the topological phase of two–dimensional gravity. Nucl. Phys., B340 (1990), 281-332.

[W2] E.Witten, Algebraic geometry associated with matrix models of two dimensional gravity, Topological models in modern mathematics (Stony Brook, NY, 1991), Publish or Perish, Houston, TX 1993, 235-269.

[Z] J.-B.Zuber, On Dubrovin's topological field theories. Mod. Phys. Lett. A9 (1994) 749-760.

Moscow State University, Moscow, Russia.

Independent University of Moscow, Moscow, Russia.

e-mail: natanzon@mccme.ru